

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



33

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07D 261/04, A61K 31/42</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/14681</b> <b>(43) International Publication Date:</b> 1 June 1995 (01.06.95)
<b>(21) International Application Number:</b> PCT/IB94/00333 <b>(22) International Filing Date:</b> 26 October 1994 (26.10.94)  <b>(30) Priority Data:</b> 08/157,248 26 November 1993 (26.11.93) US 08/262,086 17 June 1994 (17.06.94) US  <b>(60) Parent Applications or Grants</b> <b>(63) Related by Continuation</b> US 08/157,248 (CIP) Filed on 26 November 1993 (26.11.93) US 08/262,086 (CIP) Filed on 17 June 1994 (17.06.94)  <b>(71) Applicant (for all designated States except US):</b> PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US).  <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> KLEINMAN, Edward, F. [US/US]; 16 Roseridge Court, Pawcatuck, CT 06379 (US).  <b>(74) Agents:</b> SPIEGEL, Allen, J. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).		<b>(81) Designated States:</b> AU, BR, CA, CN, CZ, HU, JP, KR, NO, NZ, PL, RU, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> ISOXAZOLINE COMPOUNDS AS ANTIINFLAMMATORY AGENTS  <b>(57) Abstract</b>  This invention relates to new isoxazoline compounds which are selective inhibitors of phosphodiesterase type IV (PDE <sub>IV</sub> ). The new isoxazoline compounds are useful in inhibiting PDE <sub>IV</sub> and in the treatment of AIDS, asthma, arthritis, bronchitis, chronic obstructive pulmonary disease, psoriasis, allergic rhinitis, dermatitis, shock, atopic dermatitis, rheumatoid arthritis and osteoarthritis. This invention also relates to pharmaceutical compositions useful therefor.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

5

## ISOXAZOLINE COMPOUNDS AS ANTIINFLAMMATORY AGENTS

Background of the Invention

This invention relates to a series of 3-aryl-2-isoxazoline-5-hydroxamic acid  
10 compounds which are selective inhibitors of phosphodiesterase type IV (PDE<sub>IV</sub>) and as  
such are useful in the treatment of AIDS, asthma, arthritis, bronchitis, chronic obstructive  
pulmonary disease, psoriasis, allergic rhinitis, dermatitis and other inflammatory  
conditions.

This invention also relates to the pharmaceutically acceptable salts of said  
15 compounds; to a method of using such compounds in inhibiting PDE<sub>IV</sub>, and in the  
treatment of inflammatory conditions, AIDS, asthma, arthritis, bronchitis, chronic  
obstructive pulmonary disease, psoriasis, allergic rhinitis and dermatitis in mammals,  
especially humans; and to pharmaceutical compositions useful therefor.

The "inflammatory conditions" which can be treated according to this invention  
20 include, but are not limited to, chronic obstructive pulmonary disease, shock, atopic  
dermatitis, bronchitis, rheumatoid arthritis and osteoarthritis.

Since the recognition that cyclic AMP is an intracellular second messenger (E. W.  
Sutherland, and T. W. Rall, Pharmacol. Rev., 1960, 12, 265), inhibition of the  
phosphodiesterases has been a target for modulation and, accordingly, therapeutic  
25 intervention in a range of disease processes. More recently, distinct classes of PDE  
have been recognized (J. A. Beavo and D. H. Reifsnyder, TIPS, 1990, 11, 150), and their  
selective inhibition has led to improved drug therapy (C. D. Nicholson, R. A. Challiss  
and M. Shahid, TIPS, 1991, 12, 19). More particularly, it has been recognized that  
inhibition of PDE<sub>IV</sub> can lead to inhibition of inflammatory mediator release (M. W.  
30 Verghese et al., J. Mol. Cell Cardiol., 1989, 12 (Suppl. II), S 61) and airway smooth  
muscle relaxation (T. J. Torphy in Directions for New Anti-Asthma Drugs, eds S. R.  
O'Donnell and C. G. A. Persson, 1988, 37, Birkhauser-Verlag). Thus, compounds that  
inhibit PDE<sub>IV</sub>, but which have poor activity against other PDE types, would inhibit the  
release of inflammatory mediators and relax airway smooth muscle without causing  
35 cardiovascular effects or antiplatelet effects.

Certain pyrimidone compounds have been disclosed to be useful as antidepressants by Saccomano et al., in European Patent Application EPO 247 725 A2 published

-2-


December 2, 1987. The same pyrimidone compounds have been disclosed to be useful against asthma and certain skin disorders in International Patent Application No. PCT/US90/02162, published May 30, 1991 as International Publication Number WO 91/07178.

5

### Summary of the Invention

This invention is concerned with a series of 3-aryl-2-isoxazoline-5-hydroxamic acid compounds and to the pharmaceutically acceptable salts of such compounds. These new compounds possess inhibitory activity against PDE<sub>IV</sub> and as such are useful in treating inflammatory conditions, AIDS, asthma, arthritis, bronchitis, chronic obstructive pulmonary disease, psoriasis, allergic rhinitis or dermatitis in a mammal, especially humans.

The compounds of the present invention are of the formula (I)



the racemic, racemic-diastereomeric mixtures and optical isomers of said compounds and the pharmaceutically acceptable salts thereof wherein

m is 0, 1, 2 or 3;

n is 0, 1, 2 or 3;

25 Y<sup>1</sup> and Y<sup>2</sup> are independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, optionally substituted phenylalkyl having 1 to 6 carbons in the alkyl portion, optionally substituted phenoxyalkyl having 1 to 6 carbons in the alkyl portion, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, difluoromethyl, trifluoromethyl, fluoro, chloro, bromo, iodo, -OR<sup>1</sup> and -OR<sup>2</sup>;

30 wherein the aromatic portion of the optionally substituted phenylalkyl, and the aromatic portion of the optionally substituted phenoxyalkyl are optionally independently substituted with (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, halogen or CF<sub>3</sub>;

R<sup>1</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl, phenylalkyl having one to four carbon atoms in the alkyl portion fluoromethyl, difluoromethyl, trifluoromethyl, or -(CH<sub>2</sub>)<sub>q</sub>-quinoline wherein q is 1, 2

-3-

or 3;

$R^2$  is  $(C_1-C_3)$ alkyl,  $(C_3-C_7)$ cycloalkyl, alkoxyalkyl having 3 to 7 carbons in the alkoxy portion and 2 to 4 carbons in the alkyl portion, optionally substituted phenoxyalkyl having 2 to 6 carbons in the alkyl portion, optionally substituted phenylalkyl having 1 to 6 carbons in the alkyl portion, bicycloalkyl having 6 to 9 carbons or optionally substituted indanyl;

wherein the aromatic portion of the optionally substituted phenylalkyl, the aromatic portion of the optionally substituted phenoxyalkyl and the optionally substituted indanyl are optionally substituted with  $(C_1-C_4)$ alkyl,  $(C_1-C_4)$ alkoxy, halogen or  $CF_3$ ;

$R^3$  is hydrogen,  $(C_1-C_3)$ alkyl, fluoro $(C_1-C_3)$ alkyl having 1 to 3 fluoro atoms, mono-hydroxyalkyl having 1 to 3 carbons or alkoxyalkyl having 1 to 3 carbons in the alkyl portion and 1 to 3 carbons in the alkoxy portion;

$R^4$  is hydrogen,  $(C_1-C_3)$ alkyl, fluoro $(C_1-C_3)$ alkyl having 1 to 3 fluoro atoms, mono-hydroxyalkyl having 1 to 3 carbons, phenyl, alkoxyalkyl having 1 to 3 carbons in the alkyl portion and 1 to 3 carbons in the alkoxy portion, aminoalkyl having 1 to 3

carbons,  $-(CH_2)_n-NH-C(=O)X^1$  wherein  $X^1$  is  $(C_1-C_3)$ alkyl and n is an integer from 1

to 3, N-alkylaminoalkyl having 1 to 3 carbons in the alkylamino portion and 1 to 3 carbons in the alkyl portion,  $(C_3-C_7)$ cycloalkyl or N,N-dialkylaminoalkyl having a total of 2 to 6 carbons in the dialkylamino portion and 1 to 3 carbons in the alkyl portion;  $R^5$  is hydrogen or  $(C_1-C_3)$ alkyl;

or  $R^3$  and  $R^4$  are taken together with the carbon atoms to which they are attached and form a carbocyclic ring having 4 to 7 carbon atoms.

A preferred group of compounds or the pharmaceutically acceptable salts thereof are those compounds of the formula (I) wherein  $Y^1$  is  $-OR^1$  and is attached to the 4-position of the phenyl ring;  $Y^2$  is  $-OR^2$  and is attached to the 3-position of the phenyl ring and m, n,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  are as defined above for formula (I).

A more preferred group of compounds or the pharmaceutically acceptable salts thereof are those compounds of the formula (I) wherein  $Y^1$  is  $-OR^1$  and is attached to the 4-position of the phenyl ring and  $Y^2$  is  $-OR^2$  and is attached to the 3-position of the

-4-

phenyl ring wherein  $R^1$  is  $(C_1-C_4)$ alkyl, phenylalkyl having one to four carbon atoms in the alkyl portion or  $-(CH_2)_q$ -quinoline;  $m$  is 0;  $n$  is 0; and  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  are as defined above for formula (I).

Another more preferred group of compounds or the pharmaceutically acceptable salts thereof are those compounds of the formula (I) wherein  $Y^1$  is  $-OR^1$  and is attached to the 4-position of the phenyl ring and  $Y^2$  is  $-OR^2$  and is attached to the 3-position of the phenyl ring wherein  $R^1$  is  $(C_1-C_4)$ alkyl, phenylalkyl having one to four carbon atoms in the alkyl portion or  $-(CH_2)_q$ -quinoline;  $R^2$  is phenylalkyl having 1 to 6 carbons in the alkyl portion,  $(C_3-C_7)$ cycloalkyl, or  $(C_1-C_3)$ alkyl;  $m$  is 0;  $n$  is 0; and  $R^3$ ,  $R^4$  and  $R^5$  are as defined above for formula (I).

An even more preferred group of compounds or the pharmaceutically acceptable salts thereof are those compounds of the formula (I) wherein  $Y^1$  is  $-OR^1$  and is attached to the 4-position of the phenyl ring and  $Y^2$  is  $-OR^2$  and is attached to the 3-position of the phenyl ring wherein  $R^1$  is  $(C_1-C_4)$ alkyl, phenylalkyl having one to four carbon atoms in the alkyl portion or  $-(CH_2)_q$ -quinoline;  $R^2$  is 5-phenylpentyl, benzyl, cyclopentyl or methyl;  $m$  is 0;  $n$  is 0; and  $R^3$ ,  $R^4$  and  $R^5$  are as defined above for formula (I).

A particularly preferred group of compounds or the pharmaceutically acceptable salts thereof are those compounds of the formula (I) wherein  $Y^1$  is  $-OR^1$  and is attached to the 4-position of the phenyl ring and  $Y^2$  is  $-OR^2$  and is attached to the 3-position of the phenyl ring wherein  $R^1$  is  $(C_1-C_4)$ alkyl, phenylalkyl having one to four carbon atoms in the alkyl portion or  $-(CH_2)_q$ -quinoline;  $R^2$  is 5-phenylpentyl, benzyl, cyclopentyl or methyl;  $m$  is 0;  $n$  is 0;  $R^3$  is hydrogen and  $R^4$  and  $R^5$  are as defined above for formula (I).

Another particularly preferred group of compounds or the pharmaceutically acceptable salts thereof are those compounds of the formula (I) wherein  $Y^1$  is  $-OR^1$  and is attached to the 4-position of the phenyl ring and  $Y^2$  is  $-OR^2$  and is attached to the 3-position of the phenyl ring wherein  $R^1$  is  $(C_1-C_4)$ alkyl, phenylalkyl having one to four carbon atoms in the alkyl portion or  $-(CH_2)_q$ -quinoline;  $R^2$  is 5-phenylpentyl, benzyl, cyclopentyl or methyl;  $m$  is 0;  $n$  is 0;  $R^3$  is hydrogen;  $R^4$  is hydrogen or  $(C_1-C_5)$ alkyl and  $R^5$  is as defined above for formula (I).

A more particularly preferred group of compounds or the pharmaceutically acceptable salts thereof are those compounds of the formula (I) wherein  $Y^1$  is  $-OR^1$  and is attached to the 4-position of the phenyl ring and  $Y^2$  is  $-OR^2$  and is attached to the 3-



position of the phenyl ring wherein  $R^1$  is  $(C_1-C_4)$ alkyl, phenylalkyl having one to four carbon atoms in the alkyl portion or  $-(CH_2)_q$ -quinoline;  $R^2$  is 5-phenylpentyl, benzyl, cyclopentyl or methyl;  $m$  is 0;  $n$  is 0;  $R^3$  is hydrogen;  $R^4$  is hydrogen or  $(C_1-C_5)$ alkyl and  $R^5$  is hydrogen or  $(C_1-C_3)$ alkyl; a compound or the pharmaceutically acceptable salt thereof of the formula (I) wherein  $Y^1$  is  $-OR^1$  and is attached to the 4-position of the phenyl ring and  $Y^2$  is  $-OR^2$  and is attached to the 3-position of the phenyl ring wherein  $R^1$  is methyl;  $R^2$  is cyclopentyl;  $m$  is 0;  $n$  is 0;  $R^3$  is hydrogen;  $R^4$  is hydrogen; and  $R^5$  is hydrogen; and the levorotatory (negative rotation) isomer of a compound or the pharmaceutically acceptable salt thereof of the formula (I) wherein  $Y^1$  is  $-OR^1$  and is attached to the 4-position of the phenyl ring and  $Y^2$  is  $-OR^2$  and is attached to the 3-position of the phenyl ring wherein  $R^1$  is methyl;  $R^2$  is cyclopentyl;  $m$  is 0;  $n$  is 0;  $R^3$  is hydrogen;  $R^4$  is hydrogen; and  $R^5$  is hydrogen.

Another more particularly preferred group of compounds or the pharmaceutically acceptable salts thereof are those compounds of the formula (I) wherein  $Y^1$  is  $-OR^1$  and is attached to the 4-position of the phenyl ring and  $Y^2$  is  $-OR^2$  and is attached to the 3-position of the phenyl ring wherein  $R^1$  is methyl;  $R^2$  is cyclopentyl;  $m$  is 0;  $n$  is 0;  $R^3$  is hydrogen;  $R^4$  is methyl; and  $R^5$  is hydrogen; and the levorotatory (negative rotation) isomer of a compound or the pharmaceutically acceptable salt thereof of the formula (I) wherein  $Y^1$  is  $-OR^1$  and is attached to the 4-position of the phenyl ring and  $Y^2$  is  $-OR^2$  and is attached to the 3-position of the phenyl ring wherein  $R^1$  is methyl;  $R^2$  is cyclopentyl;  $m$  is 0;  $n$  is 0;  $R^3$  is hydrogen;  $R^4$  is methyl; and  $R^5$  is hydrogen.

The term alkyl encompasses both straight and branched chains. The aromatic portion of the optionally substituted phenylalkyl, the aromatic portion of the optionally substituted phenoxyalkyl and the optionally substituted indanyl may be substituted by one or more substituents.


#### Detailed Description of the Invention

The compounds of the present invention having the formula (I) are comprised of the racemic, racemic-diastereomeric mixtures and optical isomers of said compounds and the pharmaceutically acceptable salts thereof. The compounds of the present invention, having the formula I as defined above, are readily and generally prepared by the following reaction process.

To an alcoholic solution of sodium methoxide is added an alcoholic solution of


-6-

hydroxylamine hydrochloride and a compound of the formula



wherein  $Y^1$ ,  $Y^2$ ,  $R^3$ ,  $R^4$ ,  $m$  and  $n$  are as defined above for the compound of formula (I), and  $X$  is an alkyl group. The reaction mixture is stirred for about 12 to 24 hours, preferably 16 hours, at room temperature. The solvent is evaporated and the residue  
 5 is worked-up according to methods well known to those skilled in the art.

The intermediate ester compounds of the formula




wherein  $Y^1$ ,  $Y^2$ ,  $R^3$ ,  $R^4$ ,  $m$  and  $n$  are as defined above for the compound of formula (I), and  $X$  is an alkyl group, are synthesized according to the following procedure. To a mixture of N-chlorosuccinimide and pyridine in an inert solvent, such as methylene



-7-


chloride, is added an oxime of the formula



wherein Y<sup>1</sup> and

Y<sup>2</sup> are as defined above for formula (I). The mixture is allowed to stir for about 2 to 5

hours, preferably about 2 hours. A compound of the formula



, wherein

R<sup>3</sup> and R<sup>4</sup> are as defined above for formula I and X is an alkyl group, is added followed  
 5 by the addition of triethylamine to the mixture and the mixture stirred for about 2 hours  
 more at room temperature. The reaction is worked up according to methods well  
 known to those skilled in the art.

The synthetic method outlined above together with the following examples  
 describe methods which were and can be employed to prepare the compounds of this  
 10 invention.

Where possible, as ascertained by one skilled in the art enabled by this disclosure,  
 pharmaceutically-acceptable acid addition salts of certain compounds of this invention  
 include, but are not limited to, those formed with HCl, HBr, HNO<sub>3</sub>, H<sub>2</sub>SO<sub>4</sub>, H<sub>3</sub>PO<sub>4</sub>,  
 CH<sub>3</sub>SO<sub>3</sub>H, p-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>SO<sub>3</sub>H, CH<sub>3</sub>CO<sub>2</sub>H, gluconic acid, tartaric acid, maleic acid and  
 15 succinic acid. In the case of those compounds of the formula (I) which contain a further  
 basic nitrogen, it will, of course, be possible to form diacid addition salts (e.g., the  
 dihydrochloride) as well as the usual monoacid addition salt. Where possible, as  
 ascertained by one skilled in the art enabled by this disclosure, pharmaceutically-  
 acceptable cationic salts of certain compounds of this invention include, but are not  
 20 limited to, those of sodium, potassium, calcium, magnesium, ammonium, N,N'-dibenzyl-

-8-

ethylenediamine, N-methylglucamine (meglumine), ethanolamine and diethanolamine.

The starting materials and reagents required for the synthesis of the compounds of the present invention are readily available, either commercially, according to literature methods, or by methods exemplified in Preparations below.

- 5        The ability of the compounds or the pharmaceutically acceptable salts thereof to inhibit  $PDE_V$  and, consequently, demonstrate their effectiveness for treating inflammatory conditions is shown by the following *in vitro* assays.

#### BIOLOGICAL ASSAYS

##### Human Lung $PDE_V$

- 10        Thirty to forty grams of human lung tissue is placed in 50 mL of pH 7.4 Tris/phenyl-methylsulfonyl fluoride (PMSF)/sucrose buffer and homogenized using a Tekmar Tissumizer® (Tekmar Co., 7143 Kemper Road, Cincinnati, Ohio 45249) at full speed for 30 seconds. The homogenate is centrifuged at 48,000 x g for 70 minutes at 4°C. The supernatant is filtered twice through a 0.22  $\mu$ m filter and applied to a Mono-Q FPLC
- 15        column (Pharmacia LKB Biotechnology, 800 Centennial Avenue, Piscataway, New Jersey 08854) pre-equilibrated with pH 7.4 Tris/PMSF buffer. A flow rate of 1 mL/minute is used to apply the sample to the column, followed by a 2 mL/minute flow rate for subsequent washing and elution. Sample is eluted using an increasing, step-wise NaCl gradient in the pH 7.4 Tris/PMSF buffer. Eight mL fractions are collected. Fractions are
- 20        assayed for specific  $PDE_V$  activity, determined by [ $^3$ H]cAMP hydrolysis and the ability of a known  $PDE_V$  inhibitor (e.g. rolipram) to inhibit that hydrolysis. Appropriate fractions are pooled, diluted with ethylene glycol (2 mL ethylene glycol/5 mL of enzyme prep) and stored at -20°C until use.

- Compounds are dissolved in DMSO at a concentration of 10 mM and diluted 1:25
- 25        in water (400  $\mu$ M compound, 4% DMSO). Further serial dilutions are made in 4% DMSO to achieve desired concentrations. Final DMSO concentration in the assay tube is 1%. In duplicate the following are added, in order, to a 12 x 75 mm glass tube (all concentrations are given as final concentrations in assay tube).

- i) 25  $\mu$ l compound or DMSO (1%, for control and blank)
- 30        ii) 25  $\mu$ l pH 7.5 Tris buffer
- iii) [ $^3$ H]cAMP (1  $\mu$ M)
- iv) 25  $\mu$ l  $PDE_V$  enzyme (for blank, enzyme is preincubated in boiling water for 5 minutes)

The reaction tubes are shaken and placed in a water bath (37°C) for 20 minutes, at which time the reaction is stopped by placing the tubes in a boiling water bath for 4 minutes. Washing buffer (0.5 mL, 0.1 M 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES)/0.1 M NaCl, pH 8.5) is added to each tube on an ice bath. The contents  
5 of each tube are applied to an Affi-Gel 601 column (Biorad Laboratories, P.O. Box 1229, 85A Marcus Drive, Melville, New York 11747) (boronate affinity gel, 1 mL bed volume) previously equilibrated with washing buffer. [<sup>3</sup>H]cAMP is washed with 2 x 6 mL washing buffer, and [<sup>3</sup>H]5'AMP is then eluted with 4 mL of 0.25M acetic acid. After vortexing, 1 mL of the elution is added to 3 mL of scintillation fluid in a suitable vial, vortexed and  
10 counted for [<sup>3</sup>H].

% Inhibition is determined by the formula:

$$\% \text{ Inhibition} = 1 - \frac{\text{average cpm (test compound)} - \text{average cpm (blank)}}{\text{average cpm (control)} - \text{average cpm (blank)}} \times 100$$

IC<sub>50</sub> is defined as that concentration of compound which inhibits 50% of specific  
15 hydrolysis of [<sup>3</sup>H]cAMP to [<sup>3</sup>H]5'AMP.

For administration to humans to inhibit PDE<sub>V</sub> and in the treatment of inflammatory conditions, AIDS, asthma, arthritis, bronchitis, chronic obstructive pulmonary disease, psoriasis, allergic rhinitis or dermatitis, oral dosages of the compounds of formula (I) or the pharmaceutically acceptable salts thereof, are generally in the range of from 0.1-  
20 500 mg daily for an average adult patient (70 kg). Thus for a typical adult patient, individual tablets or capsules contain from 0.1 to 50 mg of active compound, in a suitable pharmaceutically acceptable vehicle or carrier. Multiple tablets or capsules may be required to meet the dosage requirements. Dosages for intravenous administration are typically within the range of 0.1 to 10 mg per single dose as required. For  
25 intranasal or inhaler administration, the dosage is generally formulated as a 0.1 to 1% (w/v) solution. In practice the physician will determine the actual dosage which will be most suitable for an individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case but there can, of course, be individual instances where higher or lower dosage ranges are  
30 merited, and all such dosages are within the scope of this invention.

For human use, the compounds of the formula (I) and the pharmaceutically acceptable salts thereof can be administered alone, but will generally be administered in an admixture with a pharmaceutical diluent or carrier selected with regard to the

-10-

intended route of administration and standard pharmaceutical practice. For example, they may be administered orally in the form of tablets containing such excipients as starch or lactose, or in capsules or ovals either alone or in admixture with excipients, or in the form of elixirs or suspensions containing flavoring or coloring agents. They  
5 may be injected parenterally; for example, intravenously, intramuscularly or subcutaneously. For parenteral administration, they are best used in the form of a sterile aqueous solution which may contain other substances; for example, enough salts or glucose to make the solution isotonic. For topical administration, they are best used in the form of solutions, lotions, ointments, salves and the like.

10 Thus in a further aspect the invention provides pharmaceutical compositions comprising a compound of the formula (I), or a pharmaceutically acceptable salts thereof, together with a pharmaceutically acceptable diluent or carrier which are useful: in inhibiting  $PDE_{IV}$ ; in the treatment of inflammatory conditions and in the treatment of AIDS, asthma, arthritis, bronchitis, chronic obstructive pulmonary disease, psoriasis,  
15 allergic rhinitis and dermatitis in mammals, especially humans.

This invention also provides methods of inhibiting  $PDE_{IV}$  in a mammal in need thereof which methods comprise administering to said mammal a  $PDE_{IV}$  inhibiting amount of a compound of the formula (I) or a pharmaceutically acceptable salt thereof.

This invention further provides a method of treating an inflammatory condition in  
20 a mammal in need thereof which comprises administering to said mammal an antiinflammatory amount of a compound of the formula (I) or a pharmaceutically acceptable salt thereof.

Further still, this invention provides a method of treating AIDS, asthma, arthritis, bronchitis, chronic obstructive pulmonary disease, psoriasis, allergic rhinitis, dermatitis  
25 or shock in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound or a pharmaceutically acceptable salt thereof.

The present invention is illustrated by the following examples, but it is not limited to the details thereof.

#### EXAMPLE 1

30 3-(3-Cyclopentyloxy-4-methoxy)phenyl-2-isoxazoline-5-hydroxamic Acid


To a solution of sodium methoxide, prepared from 97 mg (4.2 mmol) of sodium and 10 mL of methanol, was added 146 mg (2.1 mmol) of hydroxylamine hydrochloride in a solution of 3 mL of methanol followed by 500 mg (1.5 mmol) of the compound of

-11-

Preparation 10. After stirring for about 16 h at RT, the solvent was evaporated and the residue was dissolved in 50 mL of water and washed with ether (2 x 50 mL). The aqueous layer was acidified to pH 1 with aqueous HCl solution and the precipitate (231 mg) was filtered and recrystallized twice from CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to give 52 mg of the title compound, mp 167-168°C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 1.54-1.92 (8H, m), 3.48-3.67 (2H, m), 3.78 (3H, s), 4.79-4.85 (1H, m), 4.95 (1H, t, J=8), 6.99 (1H, d, J=9), 7.17 (1H, d, J=9), 7.23 (1H, s), 9.03 (1H, s); Anal. Calc'd. for C<sub>16</sub>H<sub>20</sub>N<sub>2</sub>O<sub>5</sub>: C, 59.99; H, 6.29; N, 8.74. Found: C, 59.82; H, 6.05; N, 8.65.

EXAMPLES 2 - 16

10 The following compounds having the formula shown below were prepared substantially according to the procedure of Example 1 substituting the indicated ester for that of the ester of Preparation 10. In the case of Example 5, N-methylhydroxylamine hydrochloride was substituted for hydroxylamine hydrochloride.






15







20

Ex	Y <sup>1</sup>	Y <sup>2</sup>	R <sup>4</sup>	R <sup>5</sup>	Ester	M.P. (°C)	Data
2	-OMe	-O(CH <sub>2</sub> ) <sub>5</sub> Ph	H	H	Cmpd. of Prep 8	130-132	<u>Anal.</u> Calc'd for C <sub>22</sub> H <sub>26</sub> N <sub>2</sub> O <sub>5</sub> : C, 66.32; H, 6.58; N, 7.03. Found: C, 66.23; H, 6.50; N, 6.94
3	-OMe	-O(CH <sub>2</sub> ) <sub>5</sub> Ph	Et	H	Cmpd. of Prep. 9	169-171	<u>Anal.</u> Calc'd for C <sub>24</sub> H <sub>30</sub> N <sub>2</sub> O <sub>5</sub> •¼H <sub>2</sub> O: C, 66.82; H, 7.07; N, 6.49. Found: C, 67.13; H, 7.03; N, 6.15

5

Ex	Y <sup>1</sup>	Y <sup>2</sup>	R <sup>4</sup>	R <sup>5</sup>	Ester	M.P. (°C)	Data
4	-OMe		Me	H	Cmpd. of Prep. 12	171-173	<i>Anal.</i> Calc'd for C <sub>17</sub> H <sub>22</sub> N <sub>2</sub> O <sub>5</sub> ·½H <sub>2</sub> O: C, 59.41; H, 6.70; N, 8.15. Found: C, 59.78; H, 6.38; N, 8.27
5	-OMe		H	Me	Cmpd. of Prep. 11	146-148	<i>Anal.</i> Calc'd for C <sub>17</sub> H <sub>22</sub> N <sub>2</sub> O <sub>5</sub> : C, 61.07; H, 6.63; N, 8.38. Found: C, 60.87; H, 6.52; N, 8.45
6	-OMe	-OMe	H	H	Cmpd. of Prep. 22	180-182	<i>Anal.</i> Calc'd for C <sub>12</sub> H <sub>14</sub> N <sub>2</sub> O <sub>5</sub> : C, 54.13; H, 5.30; N, 10.52. Found: C, 54.03; H, 5.12; N, 10.60
7	-OMe	-OCH <sub>2</sub> Ph	H	H	Cmpd. of Prep. 23	166-168	<i>Anal.</i> Calc'd for C <sub>18</sub> H <sub>18</sub> N <sub>2</sub> O <sub>5</sub> : C, 63.15; H, 5.30; N, 8.18. Found: C, 63.32; H, 5.37; N, 8.09
8		H	H	H	Cmpd. of Prep. 24	-	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> ): δ 3.48-3.68 (2H, m), 4.95 (1H, t, J=8), 5.43 (2H, s), 7.15 (2H, d, J=9), 7.59-7.69 (4H, m), 7.80 (1H, t, J=8), 8.01 (2H, t, J=7), 8.42 (1H, d, J=9), 9.04 (1H, s), 10.99 (1H, s). MS (m/e): 363 (M <sup>+</sup> )
9	-OCH <sub>2</sub> Ph	H	H	H	Cmpd. of Prep. 25	190-192	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> ): δ 3.44-3.63 (2H, m), 4.94 (1H, t, J=8), 5.15 (2H, s), 7.08 (2H, d, J=8), 7.32-7.46 (5H, m), 7.62 (2H, d, J=8), 9.03 (1H, s); MS (m/e): 313 (M <sup>+</sup> +1)


5

Ex	Y <sup>1</sup>	Y <sup>2</sup>	R <sup>4</sup>	R <sup>5</sup>	Ester	M.P. (°C)	Data
10	H		H	H	Cmpd. of Prep. 13	151-153	<i>Anal.</i> Calc'd for C <sub>15</sub> H <sub>18</sub> N <sub>2</sub> O <sub>4</sub> : C, 62.06; H, 6.25; N, 9.65. Found: C, 62.00; H, 6.15; N, 9.36
11		-OMe	H	H	Cmpd. of Prep. 14	136-138	<i>Anal.</i> Calc'd for C <sub>16</sub> H <sub>20</sub> N <sub>2</sub> O <sub>5</sub> : C, 59.99; H, 6.29; N, 8.74. Found: C, 59.66; H, 6.21; N, 8.69
12	H	H	H	H	Cmpd. of Prep. 17	166-168	<i>Anal.</i> Calc'd for C <sub>10</sub> H <sub>10</sub> N <sub>2</sub> O <sub>3</sub> : C, 58.25; H, 4.89; N, 13.59. Found: C, 58.24; H, 4.49; N, 13.45
13	OMe		Pr	H	Cmpd. of Prep. 18	154-157	<i>Anal.</i> Calc'd for C <sub>19</sub> H <sub>26</sub> N <sub>2</sub> O <sub>5</sub> : C, 62.97; H, 7.23; N, 7.73. Found: C, 62.61; H, 7.19; N, 7.54.
14	OMe		Bu	H	Cmpd. of Prep. 19	135-138	HRMS. Calc'd for C <sub>20</sub> H <sub>28</sub> N <sub>2</sub> O <sub>5</sub> : 376.19982. Found: 376.20104.
15	OMe		Ph	H	Cmpd. of Prep. 20	180-182	<i>Anal.</i> Calc'd for C <sub>22</sub> H <sub>24</sub> N <sub>2</sub> O <sub>5</sub> : C, 66.65; H, 6.10; N, 7.07. Found: C, 66.32; H, 6.30; N, 7.12.
16					Cmpd. of Prep. 21	111-133	HRMS: Calc'd for C <sub>19</sub> H <sub>24</sub> N <sub>2</sub> O <sub>5</sub> : 360.1685; Found: 360.1684.



EXAMPLES 17-22

The following compounds having the formula shown below were prepared substantially according to the procedure of Example 1 substituting the indicated ester for that of the ester of Preparation 10.



Example	Ester Cmpd. of Prep. #	R <sup>3</sup>	R <sup>4</sup>	m	MP (°C)	[α] <sub>D</sub> <sup>25</sup>	Anal.
17	28	H	H	0	137-140°	+77° MeOH	Calc'd. for C <sub>16</sub> H <sub>20</sub> N <sub>2</sub> O <sub>5</sub> : C, 59.99; H, 6.29; N, 8.74. Found: C, 59.98; H, 6.62; N, 8.80.
18	29	H	H	0	138-140°	-82° MeOH	Calc'd. for C <sub>16</sub> H <sub>20</sub> N <sub>2</sub> O <sub>5</sub> : C, 59.99; H, 6.29; N, 8.74. Found: C, 59.66; H, 6.44; N, 8.61.

Example	Ester Cmpd. of Prep. #	R <sup>3</sup>	R <sup>4</sup>	m	MP (°C)	$[\alpha]_D^{25}$	Anal.
19	33	H	H	1	139-141°	racemic	Calc'd. for C <sub>17</sub> H <sub>22</sub> N <sub>2</sub> O <sub>5</sub> ·½H <sub>2</sub> O: C, 60.02; H, 6.64; N, 8.26. Found: C, 59.63; H, 6.48; N, 8.15.
20 <sup>a</sup>	31	Me	H	0	184-186°	racemic	Calc'd. for C <sub>17</sub> H <sub>22</sub> N <sub>2</sub> O <sub>5</sub> : C, 61.01; H, 6.58; N, 8.38. Found: C, 61.08; H, 6.88; N, 8.04.
21	39	H	Me	0	167-168°	+9° <sup>b</sup> CHCl <sub>3</sub>	Calc'd. for C <sub>17</sub> H <sub>22</sub> N <sub>2</sub> O <sub>5</sub> ·½H <sub>2</sub> O: C, 60.25; H, 6.69; N, 8.27. Found: C, 60.43; H, 6.70; N, 8.23.
22	40	H	Me	0	153-155°	-14° <sup>b</sup> CHCl <sub>3</sub>	Calc'd. for C <sub>17</sub> H <sub>22</sub> N <sub>2</sub> O <sub>5</sub> ·½H <sub>2</sub> O: C, 59.41; H, 6.70; N, 8.15. Found: C, 59.64; H, 6.65; N, 8.03.

<sup>a</sup> trans isomer

<sup>b</sup> These compounds are the resolved enantiomers of Example 4. The enantiomeric purity was 99% as determined by chiral HPLC using a Chrom Tech chiral column. Mobile phase: 98:2 10 mM ammonium acetate buffer, pH 4.1: 2-propanol; flow rate: 1 mL/min; detection: 230 nm; temperature: ambient; injection volume: 20 µL.

-16-

PREPARATION 14-Methoxy-3-(5-phenylpentyl)oxy)benzaldehyde Oxime

A mixture of 25.0 g (0.164 mol) of isovanillin, 26.9 g (0.164 mol) of 5-phenyl-1-pentanol, 64.5 g (0.246 mol) of triphenylphosphine and 250 mL of THF was treated  
5 dropwise with 42.8 g (0.246 mol) of diethyl azodicarboxylate. The mixture was heated to about 90°C for about 6 hrs and then stirred overnight at RT. The solvent was evaporated and the residue was diluted with 500 mL of EtOAc, washed with water (1 x 400 mL), 1N NaOH solution (2 x 400 mL), brine (1 x 400 mL), dried (MgSO<sub>4</sub>), and evaporated to 119 g of a brown oil. Purification by flash chromatography (750 g of  
10 silica gel) using an EtOAc-hexane (3:7) eluant afforded 29.8 g (61%) of an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.42–1.92 (6H, m), 2.61 (2H, t, J=7), 3.91 (3H, s), 4.03 (2H, t, J=7), 6.91 (1H, d, J=8), 7.10–7.40 (m, 7H), 9.77 (s, 1H).


To a solution of 29.8 g (0.100 mol) of the above aldehyde in 300 mL of 95% ethanol was added 13.7 g (0.197 mol) of hydroxylamine hydrochloride in 100 mL of  
15 water followed by 16.6 g (0.197 mol) of sodium bicarbonate in small portions (gas evolution!). The mixture was stirred for about 4 h at RT and the ethanol was removed by evaporation. The residue was diluted with 250 mL of water and extracted with EtOAc (2 x 200 mL). The combined extracts were dried (MgSO<sub>4</sub>) and evaporated to a yellow oil which was crystallized from hexane/ether to afford 15.0 g of the title compound, mp  
20 65–67°C. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.46–1.93 (6H, m), 2.62 (2H, t, J=7), 3.88 (3H, s), 4.02 (2H, t, J=7), 6.99–7.62 (m, 6H), 7.49 (1H, s), 8.04 (1H, s).

An additional 2.00 g of product was obtained as a second crop from the filtrate, mp 67–69°C. Evaporation of the filtrate and purification of the residue by flash chromatography using an EtOAc-hexane (2:3) eluant also provided an additional 4.18  
25 g of product, mp 64–66°C.

PREPARATIONS 2 – 4

The following compounds having the formula shown below were prepared, substantially according to the procedure of Preparation 1, substituting the indicated phenol for isovanillin and the indicated alcohol for 5-phenyl-1-pentanol. Compounds  
30 that were oils were purified by flash chromatography.


-17-



Prep #	Y <sup>1</sup>	Y <sup>2</sup>	Phenol	Alcohol	M.P.(°C)	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ:
2	-OMe		Isovanillin	cyclopentanol	oil	1.50-2.02 (8H, m), 3.94 (3H, s), 4.62-4.80 (1H, m), 6.91 (1H, d, J=8), 6.97 (1H, dd, J=8 and 1), 7.17 (1H, d, J=1), 8.02 (1H, s), 8.16 (1H, s)
3	H		<i>m</i> -hydroxy-benzaldehyde	cyclopentanol	oil	1.50-1.95 (8H, m), 4.70-4.78 (1H, m), 6.88 (1H, dd, J=3, 8), 7.05-7.28 (3H, m), 8.09 (1H, s), 8.43 (1H, s)
4		-OMe	vanillin	cyclopentanol	110-111	1.55-2.02 (8H, m), 3.88 (3H, s), 4.78-4.88 (1H, m), 6.86 (1H, d, J=8), 7.01 (1H, dd, J=2, 8), 7.21 (1H, d, J=2), 7.65 (1H, s), 8.07 (1H, s)

PREPARATIONS 5 - 6

The following compounds having the formula shown below were prepared by condensation of the indicated aldehyde with hydroxylamine hydrochloride, substantially according to the procedure of Preparation 1.



-18-

Prep #	Y <sup>1</sup>	Y <sup>2</sup>	Aldehyde	M.P. (°C)	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> ) δ:
5	-OMe	-OH	isovanillin	146-148	3.77 (3H, s), 6.92 (2H, s), 7.08 (1H, s), 7.96 (1H, s), 9.16 (1H, s), 10.90 (1H, s)
6	-OH	H	<i>p</i> -hydroxy-benzaldehyde	115-118	6.77 (2H, d, J=9), 7.40 (2H, d, J=9), 8.00 (1H, s), 9.74 (1H, s), 10.83 (1H, s)

PREPARATION 7

10

Ethyl 2-Methylenebutyrate


A mixture of 5.0 g (0.019 mol) of triethyl 2-phosphonobutyrate, 5.5 g (0.039 mol) of K<sub>2</sub>CO<sub>3</sub>, 6.2 g (0.076 mol) of 37% aqueous formaldehyde solution, and 15 mL of water was heated to about 80°C for about 45 min. After cooling to RT, 75 mL of ether was added and the organic layer was separated, washed with brine (1 x 20 mL), dried (MgSO<sub>4</sub>), and filtered. The ether was carefully removed by distillation, leaving behind 2.1 g (87%) of the title compound as a clear oil which was used directly without further purification. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.01 (3H, t, J=7), 1.24 (3H, t, J=7), 2.26 (2H, q, J=7), 4.14 (2H, q, J=7), 5.45 (1H, s), 6.06 (1H, s).

PREPARATION 820 3-[4-Methoxy-3-(5-phenylpentyloxy)]phenyl-2-isoxazoline-5-carboxylic Acid Ethyl Ester





To a mixture of 1.28 g (9.57 mmol) of N-chlorosuccinimide, 200 μl of pyridine, and 200 mL of CH<sub>2</sub>Cl<sub>2</sub> was added 2.00 g (6.38 mmol) of the compound of Preparation 1 in a solution of 15 mL of CH<sub>2</sub>Cl<sub>2</sub>. An exotherm was observed after about 10 min and following about 2 h of stirring at RT, 644 mg (698 μl, 6.38 mmol) of ethyl acrylate was added followed by 966 mg (1.33 ml, 9.57 mmol) of triethylamine. After the exotherm subsided, the mixture was stirred for about 2 h at RT. The mixture was diluted with 250 mL of CH<sub>2</sub>Cl<sub>2</sub> and washed with aqueous 1N HCl solution, sat'd. aqueous NaHCO<sub>3</sub> solution, dried (Na<sub>2</sub>SO<sub>4</sub>), and evaporated to an oil. Purification by flash chromatography (100 g of a silica gel) using an EtOAc-hexane (2:3) eluant afforded 1.82 g (69%) of the title compound as an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.29 (3H, t, J=7), 1.40-1.91 (6H, m), 2.60 (2H, t, J=7), 3.55-3.58 (2H, m), 3.95 (3H, s), 3.99 (2H, t, J=7), 4.22 (2H, q, J=7), 5.05-5.12 (1H, m), 6.79 (1H, d, J=8), 6.95-7.31 (7H, m).

PREPARATIONS 9-21





The following compounds having the formula shown below were prepared, substantially according to the procedure of Preparation 8, substituting the indicated oxime for that of the oxime of Preparation 1 and the indicated olefin for ethyl acrylate.



Prep #	Y <sup>1</sup>	Y <sup>2</sup>	R <sup>4</sup>	X	Oxime	Olefin	M.P. (°C)	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ or Elemental Analysis:
9	-OMe	O(CH <sub>2</sub> ) <sub>5</sub> -Ph	Et	Et	Compound of Preparation 1	Compound of Preparation 7	oil	0.98 (3H, t, J=7), 1.30 (3H, t, J=7), 1.48-2.03 (8H, m), 2.62 (2H, t, J=7), 3.18 (1H, d, J=17), 3.76 (1H, d, J=17), 3.86 (3H, s), 4.01 (2H, q, J=7), 6.82 (1H, d, J=8), 7.01 (1H, dd, J=2, 8), 7.15-7.33 (6H, m)
10	-OMe		H	Et	Compound of Preparation 2	Ethyl acrylate	oil	1.27 (3H, t, J=7), 1.45-2.00 (8H, m), 3.56 (2H, d, J=10), 3.82 (3H, s), 4.22 (2H, q, J=7), 4.72-4.80 (1H, m), 5.09 (1H, t, J=10), 6.78 (1H, d, J=8), 6.98 (1H, d, J=8), 7.32 (1H, s)

Prep #	Y <sup>1</sup>	Y <sup>2</sup>	R <sup>4</sup>	X	Oxime	Olefin	M.P. (°C)	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ or Elemental Analysis:
11	-OMe		H	Me	Compound of Preparation 2	Methyl acrylate	101-102	1.45-2.00 (8H, m), 3.56 (2H, d, J=10), 3.82 (3H, s), 3.87 (3H, s), 5.16 (1H, t, J=10), 6.82 (1H, d, J=8), 7.03 (1H, dd, J=2, 8), 7.37 (1H, d, J=2)
12	-OMe		Me	Et	Compound of Preparation 2	Ethyl methacrylate	77-79	1.25 (3H, t, J=7), 1.50-2.00 (8H, m), 1.63 (3H, s), 3.11 (1H, d, J=17), 3.78 (1H, d, J=17), 3.91 (3H, s), 4.18 (2H, q, J=7), 4.68-4.77 (1H, m), 6.75 (1H, d, J=8), 6.92 (1H, dd, J=8, 2), 7.27 (1H, d, J=2)
13	H		H	Me	Compound of Preparation 3	Methyl acrylate	oil	1.45-1.95 (8H, m), 3.55-3.58 (2H, m), 3.85 (3H, s), 4.65-4.74 (1H, m), 5.09 (1H, t, J=9), 6.85 (1H, dd, J=2, 8), 7.07 (1H, d, J=8), 7.12-7.22 (2H, m)
14		OMe	H	Me	Compound of Preparation 4	Methyl acrylate	oil	1.45-1.95 (8H, m), 3.58-3.62 (2H, m), 3.79 (3H, s), 3.85 (3H, s), 4.65-4.74 (1H, m), 5.14 (1H, t, J=9), 6.83 (1H, d, J=9), 7.00 (1H, dd, J=2, 8), 7.34 (1H, d, J=2)
15	-OMe	-OH	H	Me	Compound of Preparation 5	Methyl acrylate	75-95	3.56-3.60 (2H, m), 3.79 (3H, s), 3.90 (3H, s), 5.12 (1H, t, J=10), 5.66 (1H, s), 6.83 (1H, d, J=9), 7.13 (1H, dd, J=2, 9), 7.24 (1H, d, J=2)*
16	-OH	H	H	Me	Compound of Preparation 6	Methyl acrylate	149-153	3.61-3.65 (2H, m), 3.83 (3H, s), 5.17 (1H, t, J=9), 5.80 (1H, bd s), 6.88 (2H, d, J=9), 7.56 (2H, d, J=9)



Prep #	Y <sup>1</sup>	Y <sup>2</sup>	R <sup>4</sup>	X	Oxime	Olefin	M.P. (°C)	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ or Elemental Analysis:
17	H	H	H	Et	Benzaldehyde oxime	Ethyl acrylate	40-41	1.50 (3H, t, J=7), 3.60-3.65 (2H, m), 4.26 (2H, q, J=7), 5.15 (1H, t, J=9), 7.37-7.42 (3H, m), 7.64-7.70 (2H, m)
18	OMe		Pr	Et	Compound of Preparation 2	Compound of Prep. 34	oil	0.94 (3H, t, J=8), 1.30 (3H, t, J=8), 1.32-1.99 (12 H, m), 3.20 (1H, d, J=17), 3.76 (1H, d, J=17), 3.85 (3H, s), 4.18-4.32 (2H, m), 4.72-4.82 (1H, m), 6.82 (1H, d, J=8), 7.00 (1H, dd, J=2 and 8), 7.34 (1H, d, J=2)
19	OMe		Bu	Et	Compound of Preparation 2	Compound of Prep. 35	53-36	0.90 (3H, t, J=8), 1.30 (3H, t, J=8), 1.22-1.98 (14H, m), 3.20 (1H, d, J=17), 3.78 (1H, d, J=17), 3.86 (3H, s), 4.20-4.29 (2H, m), 4.75-4.85 (1H, m), 6.82 (1H, d, J=8), 7.01 (1H, dd, J=2 and 8), 7.34 (1H, d, J=2)
20	OMe		Ph	Et	Compound of Preparation 2	Compound of Prep. 36	98-100	Anal. Calc'd for C <sub>24</sub> H <sub>27</sub> NO <sub>5</sub> : C, 70.39; H, 6.65; N, 3.42. Found: C, 70.30; H, 6.81; N, 3.56
21					Compound of Preparation 2	Methyl 1-cyclopentenoate	oil	1.60-2.37 (14H, m), 3.79 (3H, s), 3.86 (3H, s), 4.21 (1H, dd, J=3 and 7), 4.75-4.80 (1H, m), 6.82 (1H, d, J=9), 7.05 (1H, dd, J=2 and 9), 7.34 (1H, J=2)

\* The NMR shows a contaminant which is most likely a product resulting from chlorination of the aromatic ring.


-22-

PREPARATION 223-(3,4-Dimethoxyphenyl)-2-isoxazoline-5-carboxylic Acid Methyl Ester

To a solution of 1.5 g (6.00 mmol) of the compound of Preparation 15 in 25 mL of DMF was added 910 mg of  $K_2CO_3$  (6.60 mmol) and 0.41 mL (940 mg, 6.6 mmol) of methyl iodide. The mixture was heated to about 50°C and the progress of the reaction was monitored by TLC. Additional 0.4 mL portions of methyl iodide were added at about 1 and 2 h, respectively. After about 2 h of additional heating, the reaction was cooled, diluted with 250 mL of water, extracted with EtOAc (3 x 100 mL), dried ( $MgSO_4$ ), and evaporated to an oil. Purification by flash chromatography using an EtOAc-hexane (1:3) eluant afforded 270 mg of the title compound, mp 106-108°C.  $^1H$  NMR ( $CDCl_3$ ):  $\delta$  3.59-3.63 (2H, m), 3.79 (3H, s), 3.89 (3H, s), 5.15 (1H, t, J=8), 6.83 (1H, d, J=8), 7.03 (1H, dd, J=2, 8), 7.37 (1H, d, J=8); MS (m/e): 266 ( $M^+ + 1$ ).



PREPARATIONS 23 - 25

The following compounds having the formula shown below were prepared, substantially according to the procedure of Preparation 22, substituting the indicated phenol for that of Preparation 15 and the indicated alkylating agent for methyl iodide.



Prep #	Y <sup>1</sup>	Y <sup>2</sup>	Phenol	Alkylating Agent	M.P.(°C)	Data
23	OMe	OBn	Cmpd of Prep 15	PhCH <sub>2</sub> Br	183-185	$^1H$ NMR ( $CDCl_3$ ): $\delta$ 3.54-3.58 (2H,m), 3.79 (3H,s), 3.89 (3H,s), 5.10-5.26 (1H,m), 5.13 (2H,s), 6.86 (1H, d, J=9), 7.06 (1H, dd, J=2,8), 7.32-7.40 (6H,m); MS (m/e): 342 ( $M^+ + 1$ )

-23-

Prep #	Y <sup>1</sup>	Y <sup>2</sup>	Phenol	Alkylating Agent	M.P.(°C)	Data
24		H	Cmpd of Prep 16		112-113	<sup>1</sup> H NMR (CDCl <sub>3</sub> ): δ 3.56-3.59 (2H, m), 3.78 (3H, s), 5.12 (1H, t, J=8), 5.34 (2H, s), 7.02 (2H, d, J=9), 7.54-7.82 (6H, m), 8.06 (1H, d, J=8), 8.17 (1H, d, J=8); MS (m/e): 363 (M <sup>+</sup> +1)
25	-OCH <sub>2</sub> Ph	H	Cmpd of Prep 16	PhCH <sub>2</sub> Br	127-128	Anal. Calc'd for C <sub>18</sub> H <sub>17</sub> NO <sub>4</sub> : C, 69.43; H, 5.50; N, 4.50. Found: C, 69.18; H, 5.31; N, 4.59

5

PREPARATION 26[3aR-(3a $\alpha$ ,6 $\alpha$ ,7 $\alpha$ )]-Hexahydro-8,8-dimethyl-1-(1-oxo-2-propenyl)-3H-3a,6-methano-2,1-benzisothiazole 2,2-Dioxide

The title compound was prepared according to the method of Curran and Heffner (Curran, D. P., Heffner, T.A., *J. Org. Chem.*, 1990, 55, 4585) starting with (+)-L-2,10-camphor sultam, which was purchased from Fluka.

Into a 1L 3-neck round bottom flask fitted with reflux condenser, N<sub>2</sub> inlet, rubber septum and glass stopper was placed 4.03 g (0.084 mol) of 50% NaH dispersion, 400 mL of toluene, and 12.0 g (0.056 mol) of (+)-10,2-camphor sultam. After stirring for 1 h. at RT, 594 mg (0.006 mol) of CuCl followed by 9.10 mL (0.056 mol) of acryloyl chloride were added and stirring was continued overnight at RT. The mixture was then treated with 15 mL of water, evaporated, diluted with water (200 mL), and extracted with EtOAc (3 x 200 mL). The combined extracts were dried (MgSO<sub>4</sub>) and evaporated to a solid. Purification by flash chromatography (1 kg of silica gel) using a 3:7 EtOAc-hexane eluant afforded a white solid which was triturated with ether to provide 7.4 g of the title compound, mp 179-182°C.


PREPARATION 27[3aS-(3a $\alpha$ ,6 $\alpha$ ,7 $\alpha$ )]-Hexahydro-8,8-dimethyl-1-(1-oxo-2-propenyl)-3H-3a,6-methano-2,1-benzisothiazole 2,2-Dioxide

The title compound was prepared according to the procedure of Preparation 26, however, starting with (-)-D-2,10-camphor sultam, which was purchased from Fluka.


25

PREPARATIONS 28-31

The following compounds having the formula shown below were prepared substantially according to the procedure of Preparation 8, however, substituting the compound of Preparation 2 for the compound of Preparation 1 and the indicated olefins for ethyl acrylate.



Prep.	R <sup>3</sup>	R <sup>4</sup>	Q	Olefin	MP (°C)	[α] <sub>D</sub> <sup>25</sup> CHCl <sub>3</sub>	Data
28 <sup>a</sup>	H	H		Compound of Prep. 26	foam	+187°	<sup>1</sup> H-NMR(CDCl <sub>3</sub> ): δ 0.97 (3H, s), 1.19 (3H, s), 1.28-2.22 (15H, m), 3.41-3.95 (5H, m), 3.85 (3H, s), 4.74-4.82 (1H, m), 5.63 (1H, dd, J=8, 10), 6.80 (1H, d, J=8), 7.02 (1H, dd, J=2, 8), 7.33 (1H, d, J=2).

Prep.	R <sup>3</sup>	R <sup>4</sup>	Q	Olefin	MP (°C)	$[\alpha]_D^{25}$ CHCl <sub>3</sub>	Data
29 <sup>a</sup>	H	H		Compound of Prep. 27	foam	+46°	<sup>1</sup> H-NMR(CDCl <sub>3</sub> ): δ 0.94 (3H, s), 1.17 (3H, s), 1.25-2.21 (15H, m), 3.39-3.93 (5H, m), 3.83 (3H, s), 4.73-4.81 (1H, m), 5.61 (1H, dd, J=8, 10), 6.79 (1H, d, J=8), 7.01 (1H, dd, J=2, 8), 7.32 (1H, d, J=2).
30	H	H	(CH <sub>2</sub> ) <sub>2</sub> OH	CH <sub>2</sub> =CH(CH <sub>2</sub> ) <sub>2</sub> OH	89-91	racemic	Anal. Calc'd. for C <sub>17</sub> H <sub>23</sub> NO <sub>4</sub> : C, 66.86; H, 7.59; N, 4.59. Found: C, 66.71; H, 7.77; N, 4.64.
31 <sup>b</sup>	H	Me	CO <sub>2</sub> Me	ethyl crotonate	oil	racemic	<sup>1</sup> H-NMR(CDCl <sub>3</sub> ): 1.40-2.05 (8H, m), 1.43 (3H, d, J=7), 3.69 (3H, s), 3.84 (3H, s), 4.01 (1H, d, J=7), 4.74-4.82 (1H, m), 4.95-5.06 (1H, m), 6.79 (1H, d, J=8), 7.05 (1H, dd, J=2, 8), 7.31 (1H, d, J=2).

5 <sup>a</sup> less polar diastereomer (R<sub>f</sub> 0.61; 1:1 ether-toluene); <sup>b</sup> trans stereochemistry

PREPARATION 323-(3-Cyclopentyloxy-4-methoxy)phenyl-2-isoxazoline-5-acetic Acid

To a solution of 1.85 g (6.06 mmol) of the compound of Preparation 30 in 50 mL of acetone chilled to about 0°C in an ice bath was added dropwise 9.70 mL (12.1 mmol) of a 1.25 M solution of Jones reagent. The ice bath was allowed to melt, and after about 4 h. of stirring an additional 2.00 mL of Jones reagent was added and stirring was continued overnight. Excess reagent was quenched by the addition of 10 mL of isopropanol, and the solids were removed by filtration. The filtrate was concentrated and the residue was taken up in 150 mL of EtOAc, washed with water (2 x 100 mL), dried (MgSO<sub>4</sub>), and evaporated to a yellow oil. Crystallization from ether-hexane gave 1.06 g of the title compound, mp 123-126°C.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.55-2.06 (8H, m), 2.66-3.59 (4H, m), 3.87 (3H, s), 4.78-4.87 (1H, m), 5.02-5.15 (1H, m), 6.84 (1H, d, J=8), 7.02 (1H, dd, J=2, 8), 7.37 (1H, d, J=2).

PREPARATION 333-(3-Cyclopentyloxy-4-methoxy)phenyl-2-isoxazoline-5-acetic Acid Methyl Ester


A solution of 530 mg of the compound of Preparation 32 in 5 mL of MeOH was saturated with HCl gas and the mixture was stirred for about 3 h. at RT protected from atmospheric moisture with a CaCl<sub>2</sub> tube. The mixture was concentrated and the residue was taken up in 50 mL of EtOAc, washed with saturated aqueous NaHCO<sub>3</sub> solution (2 x 50 mL), dried (MgSO<sub>4</sub>), and evaporated to 530 mg of an oil. Purification by flash chromatography (25 g of silica gel) using a 2:3 - EtOAc:hexane eluant gave an oil which was crystallized from hexane-ether to afford 323 mg of the title compound as a white solid, mp 78-80°C.

Anal. Calc'd. for C<sub>18</sub>H<sub>23</sub>NO<sub>5</sub>: C, 64.85; H, 6.95; N, 4.20. Found: C, 64.49; H, 7.08; N, 4.13.

PREPARATIONS 34-36

The following compounds having the formula shown below were prepared as oils substantially according to the procedure of Preparation 7 substituting the indicated ester for triethylphosphonobutyrate.

-27-



Prep. #	R <sup>4</sup>	Ester	<sup>1</sup> H-NMR(CDCl <sub>3</sub> ):δ
5 34	Pr	triethyl phosphonopentanoate	0.90 (3H, t, J=7), 1.28 (3H, t, J=7), 1.40-1.53 (2H, m), 2.25 (2H, dt, J=1 and 7), 4.17 (2H, q, J=7), 5.48 (1H, q, J=1), 6.11 (1H, t, J=11)
35	Bu	triethyl phosphonohexanoate	0.90 (3H, t, J=7), 1.29 (3H, t, J=7), 1.26-1.48 (4H, m), 2.28 (2H, t, J=7), 4.19 (2H, q, J=7), 5.49 (1H, q, J=1), 6.11 (1H, t, J=1)
36	Ph	triethyl phosphonophenylacetate	1.32 (3H, t, J=7), 4.28 (2H, q, J=7), 5.88 (1H, d, J=1), 6.34 (1H, d, J=1), 7.20-7.45 (5H, m)

PREPARATIONS 37 and 38

10 Less Polar Diastereomer of N-[(S)-α-Methylbenzyl]-3-(3-cyclo-  
pentyloxy-4-methoxy)phenyl-5-methyl-2-isoxazoline-5-carboxamide (Preparation 37)

More Polar Diastereomer of N-[(S)-α-Methylbenzyl]-3-(3-Cyclo-  
pentyloxy-4-methoxy)phenyl-5-methyl-2-isoxazoline-5-carboxamide (Preparation 38)

A solution of 5.00 g (14 mmol) of the compound of Preparation 12 in 100 mL of  
15 absolute ethanol was treated with 2.36 g (42 mmol) of KOH and the mixture was stirred  
for about 4 hr at RT. An additional equivalent of KOH was added and stirring was  
continued for about 3 days. The mixture was concentrated, diluted with water, acidified  
with aqueous 1N HCl solution, and extracted with EtOAc (2 x 100 mL). The combined  
extracts were dried (MgSO<sub>4</sub>), evaporated, and triturated with hexane-ether to give 3.46  
20 g of 3-[3-cyclopentyloxy-4-methoxy]phenyl-5-methyl-2-isoxazoline-5-carboxylic acid, mp  
153-154°.



-28-

A mixture of 3.00 g (94 mmol) of the above compound, 100 mL of benzene, and 2.46 mL (28.2 mmol) of oxalyl chloride was heated to reflux for about 3 hr. The mixture was concentrated, diluted with 100 mL of  $\text{CH}_2\text{Cl}_2$ , and treated with 2.42 mL (18.8 mmol) of S-(-)- $\alpha$ -methylbenzylamine. After stirring for about 16 hr at RT, the mixture was concentrated, diluted with 200 mL of EtOAc, washed with aqueous 1N HCl solution (2 x 100 mL), saturated aqueous  $\text{NaHCO}_3$  solution (2 x 100 mL), dried ( $\text{Na}_2\text{SO}_4$ ), and evaporated. The residual solid (5.76 g) was purified by flash chromatography over 600 g of silica gel using 15-20% ether-toluene as eluant. Following a 500 mL pre-fraction, 35 mL-fractions were collected. Fractions 59-68 were pooled and evaporated to give 630 mg of the compound of Preparation 37, mp 154-156°C;  $R_f$  0.20, 20% ether-toluene. Anal. calculated for  $\text{C}_{25}\text{H}_{30}\text{N}_2\text{O}_4$ : C, 71.06; H, 7.16; N, 6.63. Found: C, 71.13; H, 7.42; N, 6.76.

Fractions 82-104 were pooled and concentrated to 720 mg of a white solid which was triturated with hexane-ether to give 596 mg of a white solid, mp 165-167°C. Recrystallization from ether- $\text{CH}_2\text{Cl}_2$  afforded 435 mg of the compound of Preparation 38, mp 167-168°C. An additional 1.03 g of the compound of Preparation 38, mp 166-167°C, was obtained by recrystallization (ether- $\text{CH}_2\text{Cl}_2$ ) of the combined evaporated residues of the mother liquor and Fractions 69-81. Anal. calculated for  $\text{C}_{25}\text{H}_{30}\text{N}_2\text{O}_4$ : C, 71.06; H, 7.16; N, 6.63. Found: C, 70.89; H, 7.40; N, 6.77.

#### PREPARATION 39

##### (+)-3-(3-Cyclopentyloxy-4-methoxy)phenyl-5-methyl-2-isoxazoline-5-carboxylic Acid Methyl Ester

Into a flame-dried, 3-neck round-bottom flask under  $\text{N}_2$  was placed a suspension of 549 mg (3.56 mmol) of 26% KH in mineral oil. After removal of the mineral oil by 2 successive hexane washes, the bare hydride was suspended in 35 mL of THF and a solution of 750 mg (1.78 mmol) of the compound of Preparation 37 in 35 mL of dry THF was added dropwise. After the bubbling subsided, 161  $\mu\text{L}$  (2.67 mmol) of carbon disulfide was added. The mixture was stirred for about 16 hr at RT and was quenched by the addition of 6 mL of water. The THF was evaporated and the residue was diluted with saturated aqueous  $\text{NaHCO}_3$  solution and washed with EtOAc (2 x 100 mL). The aqueous layer was acidified to pH 3 with aqueous 6N HCl solution, extracted with EtOAc (2 x 100 mL), dried ( $\text{MgSO}_4$ ), and evaporated to 217 mg of an orange oil.

A solution of the above oil in 20 mL of MeOH was saturated with HCl gas and

-29-

stirred for about 16 hr at RT. The mixture was concentrated, diluted with 50 mL of EtOAc, dried ( $\text{MgSO}_4$ ), and evaporated to a yellow solid. Purification by flash chromatography over 12 g of silica gel using a 60% EtOAc-hexane eluant afforded 131 mg of the title compound after trituration in hexane-ether, mp 127-128°C.  $[\alpha]_D^{25} +100^\circ$   
5 (c = 0.64,  $\text{CHCl}_3$ ). Anal. calculated for  $\text{C}_{18}\text{H}_{23}\text{NO}_5 \cdot \frac{1}{4}\text{H}_2\text{O}$ : C, 63.99; H, 7.01; N, 4.15. Found: C, 64.03; H, 6.96; N, 4.15.

#### PREPARATION 40

##### (-)-3-(3-Cyclopentyloxy-4-methoxy)phenyl-5-methyl-2-isoxazoline-5-carboxamide


The title compound was prepared substantially according to Procedure 39  
10 substituting the compound of Preparation 38 for the compound of Preparation 37; mp 124-125°C;  $[\alpha]_D^{25} -101^\circ$  (c = 0.61,  $\text{CHCl}_3$ ). Anal. calculated for  $\text{C}_{18}\text{H}_{23}\text{NO}_5 \cdot \frac{1}{4}\text{H}_2\text{O}$ : C, 63.99; H, 7.01; N, 4.15. Found: C, 64.04; H, 7.00; N, 4.17.

-30-

CLAIMS

What is claimed is:

1. A compound of the formula



the racemic, racemic-diastereomeric mixtures and optical isomers of said compounds and the pharmaceutically acceptable salts thereof wherein

m is 0, 1, 2 or 3;

15 n is 0, 1, 2 or 3;

$Y^1$  and  $Y^2$  are independently selected from the group consisting of hydrogen,  $(C_1-C_6)$ alkyl, optionally substituted phenylalkyl having 1 to 6 carbons in the alkyl portion, optionally substituted phenoxyalkyl having 1 to 6 carbons in the alkyl portion,  $(C_3-C_7)$ cycloalkyl, difluoromethyl, trifluoromethyl, fluoro, chloro, bromo, iodo,  $-OR^1$  and  $-OR^2$ ;

20 wherein the aromatic portion of the optionally substituted phenylalkyl, and the aromatic portion of the optionally substituted phenoxyalkyl are optionally independently substituted with  $(C_1-C_4)$ alkyl,  $(C_1-C_4)$ alkoxy, halogen or  $CF_3$ ;

$R^1$  is  $(C_1-C_4)$ alkyl, phenylalkyl having one to four carbon atoms in the alkyl portion fluoromethyl, difluoromethyl, trifluoromethyl, or  $-(CH_2)_q$ -quinoline wherein q is 1, 2 or 3;

25  $R^2$  is  $(C_1-C_3)$ alkyl,  $(C_3-C_7)$ cycloalkyl, alkoxyalkyl having 3 to 7 carbons in the alkoxy portion and 2 to 4 carbons in the alkyl portion, optionally substituted phenoxyalkyl having 2 to 6 carbons in the alkyl portion, optionally substituted phenylalkyl having 1 to 6 carbons in the alkyl portion, bicycloalkyl having 6 to 9 carbons or optionally substituted indanyl;

30

wherein the aromatic portion of the optionally substituted phenylalkyl, the aromatic portion of the optionally substituted phenoxyalkyl and the optionally substituted indanyl are optionally

-31-

substituted with (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, halogen or CF<sub>3</sub>;

R<sup>3</sup> is hydrogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl, fluoro(C<sub>1</sub>-C<sub>3</sub>)alkyl having 1 to 3 fluoro atoms, mono-hydroxyalkyl having 1 to 3 carbons or alkoxyalkyl having 1 to 3 carbons in the alkyl portion and 1 to 3 carbons in the alkoxy portion;

- 5 R<sup>4</sup> is hydrogen, (C<sub>1</sub>-C<sub>5</sub>)alkyl, fluoro(C<sub>1</sub>-C<sub>5</sub>)alkyl having 1 to 3 fluoro atoms, mono-hydroxyalkyl having 1 to 3 carbons, phenyl, alkoxyalkyl having 1 to 3 carbons in the alkyl portion and 1 to 3 carbons in the alkoxy portion, aminoalkyl having 1 to 3

carbons,  $-(CH_2)_n-NH-C(=O)-X^1$  wherein X<sup>1</sup> is (C<sub>1</sub>-C<sub>3</sub>)alkyl and n is an integer from 1

- to 3, N-alkylaminoalkyl having 1 to 3 carbons in the alkylamino portion and 1 to 3  
10 carbons in the alkyl portion, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or N,N-dialkylaminoalkyl having a total of 2 to 6 carbons in the dialkylamino portion and 1 to 3 carbons in the alkyl portion; R<sup>5</sup> is hydrogen or (C<sub>1</sub>-C<sub>3</sub>)alkyl;

or R<sup>3</sup> and R<sup>4</sup> are taken together with the carbon atoms to which they are attached and form a carbocyclic ring having 4 to 7 carbon atoms.

- 15 2. A compound according to claim 1 or a pharmaceutically acceptable salt thereof wherein Y<sup>1</sup> is -OR<sup>1</sup> and is attached to the 4-position of the phenyl ring and Y<sup>2</sup> is -OR<sup>2</sup> and is attached to the 3-position of the phenyl ring.

3. A compound according to claim 2 or a pharmaceutically acceptable salt thereof wherein R<sup>1</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl, phenylalkyl having one to four carbon atoms in the  
20 alkyl portion or -(CH<sub>2</sub>)<sub>q</sub>-quinoline; m is 0 and n is 0.

4. A compound according to claim 3 or a pharmaceutically acceptable salt thereof wherein R<sup>2</sup> is phenylalkyl having 1 to 6 carbons in the alkyl portion, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, or (C<sub>1</sub>-C<sub>3</sub>)alkyl.

5. A compound according to claim 4 or a pharmaceutically acceptable salt  
25 thereof wherein R<sup>2</sup> is 5-phenylpentyl, benzyl, cyclopentyl or methyl.

6. A compound according to claim 5 or a pharmaceutically acceptable salt thereof wherein R<sup>3</sup> is hydrogen.

7. A compound according to claim 6 or a pharmaceutically acceptable salt thereof wherein R<sup>4</sup> is hydrogen or (C<sub>1</sub>-C<sub>5</sub>)alkyl.

- 30 8. A compound according to claim 7 or a pharmaceutically acceptable salt

thereof wherein R<sup>5</sup> is hydrogen or (C<sub>1</sub>-C<sub>3</sub>)alkyl.

9. A compound according to claim 8 or a pharmaceutically acceptable salt thereof wherein R<sup>1</sup> is methyl; R<sup>2</sup> is cyclopentyl; R<sup>4</sup> is hydrogen; and R<sup>5</sup> is hydrogen.

10. The levorotatory (negative rotation) enantiomer of the compound according to claim 9 or a pharmaceutically acceptable salt thereof.

11. A compound according to claim 8 or a pharmaceutically acceptable salt thereof wherein R<sup>1</sup> is methyl; R<sup>2</sup> is cyclopentyl; R<sup>4</sup> is methyl; and R<sup>5</sup> is hydrogen.

12. The levorotatory (negative rotation) enantiomer of the compound according to claim 11 or a pharmaceutically acceptable salt thereof.

10 13. A pharmaceutical composition comprising an effective amount of a compound according to claim 1 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable diluent or carrier.

14. A method of inhibiting phosphodiesterase type IV in a mammal in need thereof which comprises administering to said mammal a phosphodiesterase type IV  
15 inhibiting amount of a compound according to claim 1 or a pharmaceutically acceptable salt thereof.

15. A method of treating an inflammatory condition in a mammal which comprises administering to said mammal an antiinflammatory amount of a compound according to claim 1 or a pharmaceutically acceptable salt thereof.

20 16. A method of treating AIDS, asthma, arthritis, bronchitis, chronic obstructive pulmonary disease, psoriasis, allergic rhinitis, dermatitis or shock in a mammal which comprises administering to said mammal an effective amount of a compound according to claim 1 or a pharmaceutically acceptable salt thereof.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 94/00333

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 6 C07D261/04 A61K31/42

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US,A,4 731 382 (BRISTOL-MEYERS CO.) 15 March 1988 see claims 1-20 ---	1-16
Y	EP,A,0 378 991 (CIBA-GEIGY AG) 25 July 1990 see page 1, line 8 - page 1, line 12; claims 1-14,20-24 ---	1-16
P,Y	WO,A,94 02448 (THE WELLCOME FOUNDATION LIMITED) 3 February 1994 see page 8, line 8 - page 8, line 17; claims 1-15 ---	1-16
P,Y	WO,A,94 18158 (PFIZER PHARMACEUTICALS INC.) 18 August 1994 see page 8, line 3 - page 8, line 18; claims 1-11 --- -/--	1-16

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

15 December 1994

Date of mailing of the international search report

20. 01. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Herz, C

# INTERNATIONAL SEARCH REPORT

Inter- national Application No  
PCT/IB 94/00333

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO,A,92 19589 (SMITHKLINE BEECHAM CORPORATION) 12 November 1992 see page 21, line 15 - page 21, line 23; claims 1-21 ---	1-16
Y	DATABASE WPI Week 9145, Derwent Publications Ltd., London, GB; AN 91-329223 & JP,A,3 220 180 (TAIHO PHARM. K.K.) 27 September 1991 see abstract ---	1-16
Y	ARZNEIM.-FORSCH., vol.24, no.4, 1974 pages 494 - 499 U. JAHN, TH. WAGNER-JAUREGG 'Wirkungsvergleich saurer Antiphlogistika im Bradykinin-, UV-Erythem- und Rattenpfotenödem-Test' * Mi 399 * see table 5 -----	1-16



# INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter national Application No

PCT/IB 94/00333

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4731382	15-03-88	AU-B- 614949 AU-A- 8310387 DE-A- 3777986 EP-A, B 0273451 JP-A- 63264442 US-A- 4861798	19-09-91 30-06-88 07-05-92 06-07-88 01-11-88 29-08-89
EP-A-0378991	25-07-90	AU-A- 4765090 CA-A- 2007085 JP-A- 2270861 US-A- 5096919	12-07-90 05-07-90 05-11-90 17-03-92
WO-A-9402448	03-02-94	AU-B- 4717093	14-02-94
WO-A-9418158	18-08-94	NONE	
WO-A-9219589	12-11-92	US-A- 5140047	18-08-92